



Renormalization of B-meson distribution amplitudes

N. Offen*

Laboratoire de Physique Théorique
 CNRS/Univ. Paris-Sud 11 (UMR 8627),
 F-91405 Orsay, France
 E-mail: nils.offen@th.u-psud.fr

S. Descotes-Genon

Laboratoire de Physique Théorique
 CNRS/Univ. Paris-Sud 11 (UMR 8627),
 F-91405 Orsay, France
 E-mail: sebastien.descotes-genon@th.u-psud.fr

We summarize a recent calculation of the evolution kernels of the two-particle B -meson distribution amplitudes ϕ_+ and ϕ_- taking into account three-particle contributions. In addition to a few phenomenological comments, we give as a new result the evolution kernel of the combination of three-particle distribution amplitudes $\Psi_A - \Psi_V$ and confirm constraints on ϕ_+ and ϕ_- derived from the light-quark equation of motion.

International Workshop on Effective Field Theories: from the pion to the upsilon
 February 2-6 2009
 Valencia, Spain

*Speaker.

1. Introduction

Exclusive decays of B -mesons provide important tools to test the Standard Model and to search for physics beyond it. Hadronic inputs encoding soft physics are not only form factors but also light-cone distribution amplitudes (LCDAs). In particular the B -meson LCDAs enter the parametrization of the hard-scattering part of hadronic matrix elements of bilocal current operators where large momentum is transferred to the soft spectator quark [1-10]. Impressive progress has been made in the calculation of the hard scattering amplitudes entering factorization theorems, see e.g. [11-15] for the $B \rightarrow PP$ case, but one limiting factor for the extraction of fundamental parameters is the uncertainty coming from the hadronic input. Recent years have seen several analyses concerning the renormalization properties [16, 17, 18] and the shape of the B -meson LCDAs [4, 18, 19, 20, 21, 22, 23]. Up to now these analyses were restricted to the two-particle case or to leading order with the exception of [23]. Here we present the results of [24] for the renormalization of the two-particle B -meson LCDAs taking into account mixing with three-parton LCDAs and in section (2.3) the results of a new calculation for the combination of three-particle LCDAs $\Psi_A - \Psi_V$ entering the equations of motion.

2. One-loop calculation with three-parton external state

The relevant two- and three-parton distribution amplitudes are defined as B to vacuum matrix-elements of a non-local heavy-to-light operator, which reads in the two-particle case [4]

$$\langle 0 | \bar{q}_\beta(z) [z, 0] (h_v)_\alpha(0) | B(p) \rangle = -i \frac{\hat{f}_B(\mu)}{4} \left[(1 + \not{v}) \left(\tilde{\phi}_+(t) + \frac{z}{2t} [\tilde{\phi}_-(t) - \tilde{\phi}_+(t)] \right) \gamma_5 \right]_{\alpha\beta} \quad (2.1)$$

and in the three-particle case [21] (the most general decomposition without contraction with a light-like vector is given in [25]):

$$\begin{aligned} \langle 0 | \bar{q}_\beta(z) [z, uz] g G_{\mu\nu}(uz) z^\nu [uz, 0] (h_v)_\alpha(0) | B(p) \rangle \\ = \frac{\hat{f}_B(\mu) M}{4} \left[(1 + \not{v}) \left[(v_\mu z - t \gamma_\mu) (\tilde{\Psi}_A(t, u) - \tilde{\Psi}_V(t, u)) - i \sigma_{\mu\nu} z^\nu \tilde{\Psi}_V(t, u) \right. \right. \\ \left. \left. - z_\mu \tilde{X}_A(t, u) + \frac{z_\mu z}{t} \tilde{Y}_A(t, u) \right] \gamma_5 \right]_{\alpha\beta}. \end{aligned} \quad (2.2)$$

We use light-like vectors $n_\pm$ so that every vector can be decomposed as

$$\begin{aligned} q_\mu &= (n_+ \cdot q) \frac{n_{-, \mu}}{2} + (n_- \cdot q) \frac{n_{+, \mu}}{2} + q_{\perp \mu} = q_+ \frac{n_{-, \mu}}{2} + q_- \frac{n_{+, \mu}}{2} + q_{\perp \mu}, \\ n_+^2 &= n_-^2 = 0 \quad n_+ \cdot n_- = 2 \quad v = (n_+ + n_-)/2. \end{aligned} \quad (2.3)$$

The computation of the renormalisation properties of the distribution amplitudes requires us to consider matrix elements of the relevant operators

$$O_+^H(\omega) = \frac{1}{2\pi} \int dt e^{i\omega t} \langle 0 | \bar{q}(z) [z, 0] \not{n}_+ \Gamma h_v(0) | H \rangle \quad (2.4)$$

$$O_-^H(\omega) = \frac{1}{2\pi} \int dt e^{i\omega t} \langle 0 | \bar{q}(z) [z, 0] \not{n}_- \Gamma h_v(0) | H \rangle \quad (2.5)$$

$$O_3^H(\omega, \xi) = \frac{1}{(2\pi)^2} \int dt e^{i\omega t} \int du e^{i\xi u} \langle 0 | \bar{q}(z) [z, uz] g_s G_{\mu\nu}(uz) z^\nu [uz, 0] \Gamma h_v(0) | H \rangle \quad (2.6)$$




Figure 1: The three leading-order contributions to the matrix element of $O_{\pm}$ with a three-parton external state.

with z parallel to n_+ , i.e. $z_\mu = t n_{+, \mu}$, $t = v \cdot z = z_-/2$ and the path-ordered exponential in the n_+ direction:


$$[z, 0] = P \exp \left[i g_s \int_0^z dy_\mu A^\mu(y) \right] \quad (2.7)$$

$$= 1 + i g_s \int_0^1 d\alpha z_\mu A^\mu(\alpha z) - g_s^2 \int_0^1 d\alpha \int_0^\alpha d\beta z_\mu z_\nu A^\mu(\alpha z) A^\nu(\beta z) + \dots \quad (2.8)$$

The Fourier-transforms of the different distribution amplitudes are then defined via

$$\phi_{\pm}(\omega) = \frac{1}{2\pi} \int dt e^{i\omega t} \tilde{\phi}_{\pm}(t) \quad F(\omega, \xi) = \frac{1}{(2\pi)^2} \int dt \int du e^{i(\omega+u\xi)t} \tilde{F}(t, u) \quad (2.9)$$

where $F = \Psi_V, \Psi_A, X_A, Y_A$. Since the renormalization of the operators is independent of the infrared properties of the matrix-elements, we can choose an on-shell partonic external state consisting of a light quark, a heavy quark and a gluon in equation (2.6). The resulting leading-order diagrams are shown in figures 1 and 2. Next-to-leading order (NLO) diagrams are obtained by adding a gluon or a quark loop (a ghost loop) in all possible places (for a complete list of diagrams, see [24]). Since the operators give rise to δ -distributions in the $+$ -component of the momenta, we chose to proceed via the theorem of residues. To be more explicit, we decomposed the loop momentum l in light-cone components, picked up the poles in the l_- -integral and performed the $l_\perp$ -integration in dimensional regularization with $D = 2 - 2\varepsilon$ dimensions. Additional $\frac{1}{\varepsilon}$ -poles arise through the l_+ -integration for diagrams where a gluon is exchanged between the Wilson-line from the operator and the heavy-quark field. These are related to the cusp anomalous dimension, see e.g [26, 27], stemming from the intersection of one light-like Wilson line from the path ordered exponential in the operator and one time-like Wilson line from the interaction of soft gluons with the heavy quark. The additional poles give rise to $\frac{1}{\varepsilon^2}$ -terms as well as Sudakov logarithms.



$$A_{3\mu} : ig_s(q_+\varepsilon_\mu - q_\mu\varepsilon_+) \bar{v}\Gamma^a u \delta(\omega - k_+) \delta(\xi - q_+)$$

Figure 2: Leading-order contribution to the matrix element of $O_{3\mu}$ with a three-parton external state.

2.1 ϕ_+ -case

In the ϕ_+ -case there is no mixing from three-particle distribution amplitudes. $\gamma_{+,3} = 0$ at order α_s . We confirm the result for the anomalous-dimension matrix found in [16]

$$\gamma_+^{(1)}(\omega, \omega'; \mu) = \left(\Gamma_{\text{cusp}}^{(1)} \log \frac{\mu}{\omega} + \gamma^{(1)} \right) \delta(\omega - \omega') - \Gamma_{\text{cusp}}^{(1)} \omega \left(\frac{\theta(\omega' - \omega)}{\omega'(\omega' - \omega)} + \frac{\theta(\omega - \omega')}{\omega(\omega - \omega')} \right)_+ \quad (2.10)$$

with

$$\left[f(\omega, \omega') \right]_+ = f(\omega, \omega') - \delta(\omega - \omega') \int d\omega' f(\omega, \omega') \quad \Gamma_{\text{cusp}}^{(1)} = 4 \quad \gamma^{(1)} = -2$$

2.2 ϕ_- -case

The ϕ_- case is more involved. After coupling-constant and external leg renormalization there remains a genuine three-particle term. The renormalization group equation to order α_s can be written as

$$\begin{aligned} \frac{\partial \phi_-(\omega; \mu)}{\partial \log \mu} = & -\frac{\alpha_s(\mu)}{4\pi} \left(\int d\omega' \gamma_-^{(1)}(\omega, \omega'; \mu) \phi_-(\omega'; \mu) \right. \\ & \left. + \int d\omega' d\xi' \gamma_{-,3}^{(1)}(\omega, \omega', \xi'; \mu) [\Psi_A - \Psi_V](\omega', \xi'; \mu) \right) \end{aligned} \quad (2.11)$$

where $\gamma_-^{(1)}$ is the result from [17]

$$\gamma_-^{(1)}(\omega, \omega'; \mu) = \gamma_+^{(1)} - \Gamma_{\text{cusp}}^{(1)} \frac{\theta(\omega' - \omega)}{\omega'} \quad (2.12)$$

and $\gamma_{-,3}^{(1)}$ from [24]

$$\begin{aligned} \gamma_{-,3}^{(1)}(\omega, \omega', \xi') = & 4 \left[\frac{\Theta(\omega)}{\omega'} \left\{ (C_A - 2C_F) \left[\frac{1}{\xi'^2} \frac{\omega - \xi'}{\omega' + \xi' - \omega} \Theta(\xi' - \omega) + \frac{\Theta(\omega' + \xi' - \omega)}{(\omega' + \xi')^2} \right] \right. \right. \\ & \left. \left. - C_A \left[\frac{\Theta(\omega' + \xi' - \omega)}{(\omega' + \xi')^2} - \frac{1}{\xi'^2} (\Theta(\omega - \omega') - \Theta(\omega - \omega' - \xi')) \right] \right\} \right]_+ \end{aligned} \quad (2.13)$$

where we defined the $+$ -distribution with three variables as

$$\left[f(\omega, \omega', \xi') \right]_+ = f(\omega, \omega', \xi') - \delta(\omega - \omega' - \xi') \int d\omega f(\omega, \omega', \xi'') \quad (2.14)$$

2.3 $\Psi_A - \Psi_V$ -renormalization and equation of motion constraints

Here we report on an up to now unpublished calculation of the renormalization of the three-particle LCDAs $\Psi_A - \Psi_V$. We project on the relevant distribution amplitudes in equation (2.3) using $\Gamma = \gamma_\perp^\mu \not{n}_+ \not{n}_- \gamma_5$ (although a γ^μ instead of $\gamma_\perp^\mu$ yields the same result). The calculations go along the same lines as in the previous two cases, even though there is only one leading-order structure (shown in figure 2) and NLO diagrams must have one gluon attached to the vertex in order not to vanish trivially. For convenience the result is splitted into C_F - and C_A -colour structures.

$$\begin{aligned}
\gamma_{3,3,C_A}^{(1)}(\omega, \xi, \omega', \xi') = & 2 \left[\delta(\omega - \omega') \left\{ \frac{\xi}{\xi'^2} \Theta(\xi' - \xi) - \left[\frac{\Theta(\xi - \xi')}{\xi - \xi'} \right]_+ - \left[\frac{\xi}{\xi'} \frac{\Theta(\xi' - \xi)}{\xi' - \xi} \right]_+ \right\} \right. \\
& + \delta(\xi - \xi') \left\{ \left[\frac{\Theta(\omega - \omega')}{\omega - \omega'} \right]_+ + \left[\frac{\omega}{\omega'} \frac{\Theta(\omega' - \omega)}{\omega' - \omega} \right]_+ \right\} + \delta(\omega + \xi - \omega' - \xi') \\
& \times \left\{ \frac{1}{\xi'} \Theta(\omega - \omega') - \left[\frac{\Theta(\omega - \omega')}{\omega - \omega'} \right]_+ - \left[\frac{\omega}{\omega'} \frac{\Theta(\omega' - \omega)}{\omega' - \omega} \right]_+ \right\} \\
& + \delta(\omega + \xi - \omega' - \xi') \frac{1}{\xi'(\omega' + \xi')} \left\{ \frac{\omega - \xi'}{\xi'} (\omega' + \xi' - \omega) \Theta(\omega - \omega') \right. \\
& - \frac{\omega}{\omega'} (\omega' + 2\xi' - \omega) \Theta(\omega' - \omega) \Theta(\omega) + \frac{\omega}{\xi'} (\omega - \xi') \Theta(\xi' - \omega) \Theta(\omega) \\
& \left. \left. + \frac{\omega - \xi'}{\omega'} (\omega' + \xi' - \omega) \Theta(\omega - \xi') \Theta(\xi) \right\} \right] \quad (2.15)
\end{aligned}$$

$$\begin{aligned}
\gamma_{3,3,C_F}^{(1)}(\omega, \xi, \omega', \xi'; \mu) = & \gamma_+^{(1)}(\omega, \omega'; \mu) \delta(\xi - \xi') + \gamma_{R3,3}^{(1)}(\omega, \xi, \omega', \xi') \\
\gamma_{R3,3}^{(1)}(\omega, \xi, \omega', \xi') = & 4\delta(\omega + \xi - \omega' - \xi') \\
& \times \left[\frac{\xi^2}{\omega'} \frac{\Theta(\omega' - \xi)}{(\omega + \xi)^2} \Theta(\xi) + \frac{\omega}{\xi'} \frac{\Theta(\xi - \omega')}{\omega + \xi} \Theta(\omega) \left(\frac{\xi}{\omega + \xi} - \frac{\omega - \xi'}{\xi'} \right) \right] \quad (2.16)
\end{aligned}$$

with $\gamma_+^{(1)}$ the same as in equation (2.10) and $\gamma_{3,3}^{(1)}$ defined as in (2.11) with obvious changes. Part of this calculation, namely the light-quark-gluon part, has been calculated in a different context and a different scheme, e.g. in [29, 28].

In [21] two equations from the light- and heavy-quark equations of motion were derived

$$\omega \phi'_-(\omega; \mu) + \phi_+(\omega; \mu) = I(\omega; \mu), \quad (\omega - 2\bar{\Lambda}) \phi_+(\omega; \mu) + \omega \phi_-(\omega; \mu) = J(\omega; \mu), \quad (2.17)$$

where $I(J)(\omega; \mu)$ are integro-differential expressions involving the three-particle LCDAs $\Psi_A - \Psi_V$ ($\Psi_A + X_A$ and Ψ_V) respectively. While the second equation was shown not to hold beyond leading order in [17, 23] we have checked that the first one is valid once renormalization is taken into account. Taking the derivative of the first equation with respect to $\log \mu$, inserting

$$I(\omega; \mu) = 2 \frac{d}{d\omega} \int_0^\omega d\rho \int_{\omega-\rho}^\infty \frac{d\xi}{\xi} \frac{\partial}{\partial \xi} [\Psi_A(\rho, \xi; \mu) - \Psi_V(\rho, \xi; \mu)] \quad (2.18)$$

and using the relation from [17]

$$-\omega \frac{d}{d\omega} \int_0^\eta \frac{d\omega'}{\eta} \gamma_-^{(1)}(\omega, \omega'; \mu) = \gamma_+^{(1)}(\omega, \eta; \mu) \quad (2.19)$$

one arrives at the following equation

$$\begin{aligned}
& \omega \frac{d}{d\omega} \int d\omega' d\xi' \gamma_{-3}^{(1)}(\omega, \omega', \xi') (\Psi_A(\omega', \xi'; \mu) - \Psi_V(\omega', \xi'; \mu)) \\
& + 2 \int d\omega' \gamma_+^{(1)}(\omega, \omega'; \mu) \frac{d}{d\omega'} \int_0^{\omega'} d\rho \int_{\omega'-\rho}^{\infty} \frac{d\xi}{\xi} \frac{\partial}{\partial \xi} (\Psi_A(\rho, \xi; \mu) - \Psi_V(\rho, \xi; \mu)) \\
& = 2 \int d\omega' d\xi' \frac{d}{d\omega} \int_0^{\omega} d\rho \int_{\omega-\rho}^{\infty} \frac{d\xi}{\xi} \frac{\partial}{\partial \xi} \gamma_{3,3}^{(1)}(\rho, \xi, \omega' \xi'; \mu) (\Psi_A(\omega', \xi'; \mu) - \Psi_V(\omega', \xi'; \mu)),
\end{aligned} \tag{2.20}$$

which can be proven to hold at order α_s by simple insertion of the respective evolution kernels (2.10), (2.13), (2.15), (2.16). This non-trivial outcome gives us further confidence concerning the renormalization group properties of the LCDAs.

3. Conclusions

The presence of $\delta(\omega - \omega') \log(\mu/\omega)$ in the renormalization matrices gives rise to a radiative tail falling off like $(\log \omega)/\omega$ for large ω . Therefore non-negative moments of the LCDAs are not well defined and have to be considered with an ultraviolet cut-off.[16, 17, 22, 23]

$$\langle \omega^N \rangle_{\pm}(\mu) = \int_0^{\Lambda_{UV}} d\omega \omega^N \phi_{\pm}(\omega; \mu) \tag{3.1}$$

For ϕ_- it is interesting to examine the limit

$$\lim_{\Lambda_{UV} \rightarrow \infty} \int_0^{\Lambda_{UV}} d\omega \omega^N z_{-,3}^{(1)}(\omega, \omega', \xi') = 0 \quad N = 0, 1, \quad z_{-,3}^{(1)} = \frac{1}{2\varepsilon} \gamma_{-,3}^{(1)}, \tag{3.2}$$

which is relevant for the calculation of the three-particle contributions to the moments:

$$\begin{aligned}
\int_0^{\Lambda_{UV}} d\omega \omega^N \phi_-(\omega; \mu) &= 1 + \frac{\alpha_s}{4\pi} \left(\int d\omega' \phi_-(\omega') \int_0^{\Lambda_{UV}} d\omega \omega^N z_-^{(1)}(\omega, \omega'; \mu) \right. \\
&\quad \left. - \int d\omega' d\xi' (2 - D) [\Psi_A - \Psi_V](\omega', \xi') \int_0^{\Lambda_{UV}} d\omega \omega^N z_{-,3}^{(1)}(\omega, \omega', \xi') \right)
\end{aligned} \tag{3.3}$$

Therefore as stated in [17] three-particle distribution amplitudes give only subleading contribution to the first two moments ($N = 0, 1$) and we have explicitly checked that this statement cannot be extended to higher moments ($N \geq 2$).

The next step consists in using the renormalization properties as a guide to go beyond the existing models derived from a leading-order sum-rule calculation resulting in $\Psi_A = \Psi_V$ [20] and to analyze their influence on ϕ_- . Finally, for practical calculations involving three-particle contributions, one would need the evolution kernels for the all relevant LCDAs, which will be the subject of a future work.

Acknowledgments

Work supported in part by EU Contract No. MRTN-CT-2006-035482, “FLAVIANet” and by the ANR contract “DIAM” ANR-07-JCJC-0031.

References

- [1] M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, Nucl. Phys. B **591** (2000) 313 [arXiv:hep-ph/0006124].
- [2] M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, Nucl. Phys. B **606** (2001) 245 [arXiv:hep-ph/0104110].
- [3] M. Beneke and M. Neubert, Nucl. Phys. B **675** (2003) 333 [arXiv:hep-ph/0308039].
- [4] A. G. Grozin and M. Neubert, Phys. Rev. D **55**, 272 (1997) [arXiv:hep-ph/9607366].
- [5] M. Beneke and T. Feldmann, Nucl. Phys. B **685**, 249 (2004) [arXiv:hep-ph/0311335].
- [6] S. Descotes-Genon and C. T. Sachrajda, Nucl. Phys. B **650**, 356 (2003) [arXiv:hep-ph/0209216].
- [7] S. Descotes-Genon and C. T. Sachrajda, Phys. Lett. B **557**, 213 (2003) [arXiv:hep-ph/0212162].
- [8] S. Descotes-Genon and C. T. Sachrajda, Nucl. Phys. B **693**, 103 (2004) [arXiv:hep-ph/0403277].
- [9] M. Beneke, T. Feldmann and D. Seidel, Nucl. Phys. B **612**, 25 (2001) [arXiv:hep-ph/0106067].
- [10] M. Beneke and T. Feldmann, Nucl. Phys. B **592**, 3 (2001) [arXiv:hep-ph/0008255].
- [11] G. Bell, arXiv:0902.1915 [hep-ph].
- [12] G. Bell, Nucl. Phys. B **795** (2008) 1 [arXiv:0705.3127 [hep-ph]].
- [13] M. Beneke and S. Jager, Nucl. Phys. B **768** (2007) 51 [arXiv:hep-ph/0610322].
- [14] M. Beneke and S. Jager, Nucl. Phys. B **751** (2006) 160 [arXiv:hep-ph/0512351].
- [15] V. Pilipp, Nucl. Phys. B **794** (2008) 154 [arXiv:0709.3214 [hep-ph]].
- [16] B. O. Lange and M. Neubert, Phys. Rev. Lett. **91** (2003) 102001 [arXiv:hep-ph/0303082].
- [17] G. Bell and T. Feldmann, JHEP **0804** (2008) 061 [arXiv:0802.2221 [hep-ph]].
- [18] V. M. Braun, D. Y. Ivanov and G. P. Korchemsky, Phys. Rev. D **69**, 034014 (2004) [arXiv:hep-ph/0309330].
- [19] A. Khodjamirian, T. Mannel and N. Offen, Phys. Lett. B **620** (2005) 52 [arXiv:hep-ph/0504091].
- [20] A. Khodjamirian, T. Mannel and N. Offen, Phys. Rev. D **75** (2007) 054013 [arXiv:hep-ph/0611193].
- [21] H. Kawamura, J. Kodaira, C. F. Qiao and K. Tanaka, Phys. Lett. B **523** (2001) 111 [Erratum-ibid. B **536** (2002) 344] [arXiv:hep-ph/0109181].
- [22] S. J. Lee and M. Neubert, Phys. Rev. D **72** (2005) 094028 [arXiv:hep-ph/0509350].
- [23] H. Kawamura and K. Tanaka, arXiv:0810.5628 [hep-ph].
- [24] S. Descotes-Genon and N. Offen, arXiv:0903.0790 [hep-ph].
- [25] B. Geyer and O. Witzel, Phys. Rev. D **72** (2005) 034023 [arXiv:hep-ph/0502239].
- [26] G. P. Korchemsky and A. V. Radyushkin, Nucl. Phys. B **283** (1987) 342.
- [27] I. A. Korchemskaya and G. P. Korchemsky, Phys. Lett. B **287** (1992) 169.
- [28] I. I. Balitsky and V. M. Braun, Nucl. Phys. B **311** (1989) 541.
- [29] A. P. Bukhvostov, G. V. Frolov, L. N. Lipatov and E. A. Kuraev, Nucl. Phys. B **258** (1985) 601.